2007 Snake River Watershed Summary (36)

		Conservation Tillage			Total	Other Tillage Practices	
	Total	(greater than 30% residue)			= Conservation	(15-30% residue)	(0-15% residue)
Annual Crop	Points	No-Till	Ridge-Till	Mulch-Till	Tillage	Reduce-Till	Intensive-Till
Corn	5	0	0	1	1	0	4
Small Grain (Spring-Seeded)	0	0	0	0	0	0	0
Small Grain (Fall Seeded)	0	0	0	0	0	0	0
Soybeans (Full Season)	6	3	0	1	4	0	2
Soybeans (Double-Cropped)	0	0	0	0	0	0	0
Cotton	0	0	0	0	0	0	0
Grain Sorghum	0	0	0	0	0	0	0
Forage Crops	0	0	0	0	0	0	0
Other Crops	0	0	0	0	0	0	0
Total Points	11	3	0	2	5	0	6
Perm. Pasture	0						
Fallow	0						
Forages	5						

		Conservation Tillage			Total	Other Tillage Practices	
	Total	(greater than 30% residue)			= Conservation	(15-30% residue)	(0-15% residue)
Annual Crop	Points	No-Till	Ridge-Till	Mulch-Till	Tillage	Reduce-Till	Intensive-Till
Corn	5	0.0%	0.0%	20.0%	20.0%	0.0%	80.0%
Small Grain (Spring-Seeded)	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Small Grain (Fall Seeded)	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Soybeans (Full Season)	6	50.0%	0.0%	16.7%	66.7%	0.0%	33.3%
Soybeans (Double-Cropped)	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Cotton	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Grain Sorghum	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Forage Crops	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Other Crops	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Total Points	11	27.3%	0.0%	18.2%	45.5%	0.0%	54.5%

Perm. Pasture	
Fallow	
Forages	
Conservation Reserve Program	

0

0

Conservation Reserve Program